Discrete Structures for Computer Science

William Garrison
bill@cs.pitt.edu
6311 Sennott Square

Lecture #22: Primes, GCDs, and LCMs

University of Pittsburgh

Today's Topics

Primes & Greatest Common Divisors

- Prime factorizations
- Important theorems about primality
- Greatest Common Divisors
- Least Common Multiples
- Euclid's algorithm

Let's (finally) define the primes formally

Definition: A prime number is a positive integer *p* greater than 1 that is divisible by only 1 and itself. If a number is not prime, it is called a composite number.

Mathematically: p is prime $\Leftrightarrow p>1 \wedge \forall x \in \mathbf{Z}^+ [(x \neq 1 \wedge x \neq p) \rightarrow x \nmid p]$

Examples: Are the following numbers prime or composite?

- 23
- 42
- 17
- 3
- 9

Any positive integer can be represented as a unique product of prime numbers!

Theorem (The Fundamental Theorem of Arithmetic): Every positive integer greater than 1 can be written uniquely as a prime or the product of two or more primes where the prime factors are written in order of non-decreasing size.

Examples:

- $100 = 2 \times 2 \times 5 \times 5 = 2^2 \times 5^2$
- $641 = 641$
- $999 = 3 \times 3 \times 3 \times 37 = 3^3 \times 37$
- $1024 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 2^{10}$

Note: Proving the fundamental theorem of arithmetic requires some mathematical tools that we have not yet learned.

This leads to a related theorem…

Theorem: If n is a composite integer, then n has a prime divisor less than or equal to $\sqrt{n}$.

Proof:

- If n is composite, then it has a positive integer factor a with $1 < a < n$ by definition. This means that $n = ab$, where b is an integer greater than 1.
- Assume $a > \sqrt{n}$ and $b > \sqrt{n}$. Then $ab > \sqrt{n}\sqrt{n} = n$, which is a contradiction. So either $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.
- Thus, n has a divisor less than or equal to $\sqrt{n}$.
- By the fundamental theorem of arithmetic, this divisor is either prime, or is a product of primes. In either case, n has a prime divisor less than or equal to $\sqrt{n}$. ❑

Applying contraposition leads to a naive primality test

Corollary: If n is a positive integer that does not have a prime divisor less than or equal to $\sqrt{n}$, then n is prime.

Example: Is 101 prime?

- The primes less than or equal to $\sqrt{101}$ are 2, 3, 5, and 7
- Since 101 is not divisible by 2, 3, 5, or 7, it must be prime

Example: Is 1147 prime?

- The primes less than or equal to $\sqrt{1147}$ are 2, 3, 5, 7, 11, 13, 17, 23, 29, and 31
- $1147 = 31 \times 37$, so 1147 must be composite

This approach can be generalized

The Sieve of Eratosthenes is a brute-force algorithm for finding all prime numbers less than some value *n*

Step 1: List the numbers less than *n*

2	3	~~4~~	5	~~6~~	7	~~8~~	~~9~~	~~10~~	11
~~12~~	13	~~14~~	~~15~~	~~16~~	17	~~18~~	19	~~20~~	~~21~~
~~22~~	23	~~24~~	~~25~~	~~26~~	~~27~~	~~28~~	29	~~30~~	31
~~32~~	~~33~~	~~34~~	~~35~~	~~36~~	37	~~38~~	~~39~~	~~40~~	41
~~42~~	43	~~44~~	~~45~~	~~46~~	47	~~48~~	~~49~~	~~50~~	~~51~~
~~52~~	53	~~54~~	~~55~~	~~56~~	~~57~~	~~58~~	59	~~60~~	61
~~62~~	~~63~~	~~64~~	~~65~~	~~66~~	67	~~68~~	~~69~~	~~70~~	71

Step 2: If the next available number is less than √n, cross out all of its multiples

Step 3: Repeat until the next available number is > √n

Step 4: All remaining numbers are prime

How many primes are there?

Theorem: There are infinitely many prime numbers.

Proof: By contradiction

- Assume that there are only a finite number of primes $p_1, \ldots, p_n$
- Let $Q = p_1 \times p_2 \times \ldots \times p_n + 1$
- By the fundamental theorem of arithmetic, Q can be written as the product of two or more primes.
- Note that no p_j divides Q, for if $p_j \mid Q$, then p_j also divides $Q - p_1 \times p_2 \times \ldots \times p_n = 1$.
- Therefore, there must be some prime number not in our list. This prime number is either Q (if Q is prime) or a prime factor of Q (if Q is composite).
- This is a contradiction since we assumed that all primes were listed. Therefore, there are infinitely many primes. ❑

This is a non-constructive existence proof!

In-class exercises

Problem 1: What is the prime factorization of 984?

Problem 2: Is 157 prime? Is 97 prime?

Problem 3: Is the set of all prime numbers countable or uncountable? If it is countable, show a 1-to-1 correspondence between the prime numbers and the natural numbers.

Greatest common divisors

Definition: Let a and b be integers, not both zero. The largest integer d such that $d \mid a$ and $d \mid b$ is called the greatest common divisor of a and b, denoted by $\gcd(a, b)$.

Note: We can (naively) find GCDs by comparing the common divisors of two numbers.

Example: What is the GCD of 24 and 36?

- Factors of 24: 1, 2, 3, 4, 6, 8, 12, 24
- Factors of 36: 1, 2, 3, 4, 6, 9, 12, 18, 36
- $\therefore \gcd(24, 36) = 12$

Sometimes, the GCD of two numbers is 1

Example: What is gcd(17, 22)?

- Factors of 17: 1, 17
- Factors of 22: 1, 2, 11, 22
- $\therefore$ gcd(17, 22) = 1

Definition: If gcd(a, b) = 1, we say that a and b are relatively prime, or coprime. We say that a_1, a_2, …, a_n are pairwise relatively prime if gcd(a_i, a_j) = 1 $\forall i,j$.

Example: Are 10, 17, and 21 pairwise coprime?

- Factors of 10: 1, 2, 5, 10
- Factors of 17: 1, 17
- Factors of 21: 1, 3, 7, 21

We can leverage the fundamental theorem of arithmetic to develop a better algorithm

Let: $a = p_1^{a_1} p_2^{a_2} \cdots p_n^{a_n}$ and $b = p_1^{b_1} p_2^{b_2} \cdots p_n^{b_n}$

Then:

$$gcd(a, b) = p_1^{min(a_1,b_1)} p_2^{min(a_2,b_2)} \cdots p_n^{min(a_n,b_n)}$$

Greatest multiple of p_1 in both a and b

Greatest multiple of p_2 in both a and b

Example: Compute gcd(120, 500)

- $120 = 2^3 \times 3 \times 5$
- $500 = 2^2 \times 5^3$
- So gcd(120, 500) = $2^2 \times 3^0 \times 5 = 20$

Better still is Euclid's algorithm

Observation: If $a = bq + r$, then $\gcd(a, b) = \gcd(b, r)$

Proved in section 4.3 of the book

So, let $r_0 = a$ and $r_1 = b$. Then:

- $r_0 = r_1 q_1 + r_2$ $\quad 0 \le r_2 < r_1$
- $r_1 = r_2 q_2 + r_3$ $\quad 0 \le r_3 < r_2$
- …
- $r_{n-2} = r_{n-1} q_{n-1} + r_n$ $\quad 0 \le r_n < r_{n-1}$
- $r_{n-1} = r_n q_n$

$\gcd(a, b) = r_n$

Examples of Euclid's algorithm

Example: Compute gcd(414, 662)

- $662 = 414 \times 1 + 248$
- $414 = 248 \times 1 + 166$
- $248 = 166 \times 1 + 82$
- $166 = 82 \times 2 + 2$ ⟸ gcd(414, 662) = 2
- $82 = 2 \times 41$

Example: Compute gcd(9888, 6060)

- $9888 = 6060 \times 1 + 3828$
- $6060 = 3828 \times 1 + 2232$
- $3828 = 2232 \times 1 + 1596$
- $2232 = 1596 \times 1 + 636$
- $1596 = 636 \times 2 + 324$
- $636 = 324 \times 1 + 312$
- $324 = 312 \times 1 + 12$ ⟸ gcd(9888, 6060) = 12
- $312 = 12 \times 26$

Least common multiples

Definition: The least common multiple of the integers a and b, where neither is 0, is the smallest positive integer that is divisible by both a and b. The least common multiple of a and b is denoted $\text{lcm}(a, b)$.

Example: What is lcm(3,12)?

- Multiples of 3: 3, 6, 9, 12, 15, …
- Multiples of 12: 12, 24, 36, …
- So lcm(3,12) = 12

Note: $\text{lcm}(a, b)$ is guaranteed to exist, since a common multiple exists (i.e., ab).

We can leverage the fundamental theorem of arithmetic to develop a better algorithm

Let: $a = p_1^{a_1} p_2^{a_2} \cdots p_n^{a_n}$ and $b = p_1^{b_1} p_2^{b_2} \cdots p_n^{b_n}$

Then:

$$lcm(a, b) = p_1^{max(a_1,b_1)} p_2^{max(a_2,b_2)} \cdots p_n^{max(a_n,b_n)}$$

Greatest multiple of p_1 in either a or b

Greatest multiple of p_2 in either a or b

Example: Compute lcm(120, 500)

- $120 = 2^3 \times 3 \times 5$
- $500 = 2^2 \times 5^3$
- So lcm(120, 500) = $2^3 \times 3 \times 5^3$ = 3000 << 120×500 = 60,000

LCMs are closely tied to GCDs

Note: $ab = \text{lcm}(a, b) \times \gcd(a, b)$

Example: $a = 120 = 2^3 \times 3 \times 5$, $b = 500 = 2^2 \times 5^3$

- $120 = 2^3 \times 3 \times 5$
- $500 = 2^2 \times 5^3$
- $\text{lcm}(120, 500) = 2^3 \times 3 \times 5^3 = 3000$
- $\gcd(120, 500) = 2^2 \times 3^0 \times 5 = 20$
- $\text{lcm}(120, 500) \times \gcd(120, 500)$

✔

In-class exercises

Problem 4: Use Euclid's algorithm to compute gcd(92928, 123552).

Problem 5: Compute gcd(24, 36) and lcm(24, 36). Verify that gcd(24, 36) $\times$ lcm(24, 36) = 24 $\times$ 36.

(Submit both on Top Hat after a few minutes to work)

Final Thoughts

- Prime numbers play an important role in number theory
- There are an infinite number of prime numbers
- Any number can be represented as a product of prime numbers; this has implications when computing GCDs and LCMs
- Next time: Solving congruences, modular inverses