



Homework 5

- Minimum Value 63.00
- Maximum Value 100.00
- Average 88.88

- 90 - 100 22
- 80 - 89 10
- 70 - 79 6
- 60 - 69 2
- Null 7



Today's Topics

Integers and division

- The division algorithm
- Modular arithmetic
- Applications of modular arithmetic



What is number theory?

Number theory is the branch of mathematics that explores the integers and their properties.

Number theory has many applications within computer science, including:

- Organizing data
- Encrypting sensitive data
- Developing error correcting codes
- Generating “random” numbers
- ...

We will only scratch the surface...



The notion of divisibility is one of the most basic properties of the integers

Definition: If a and b are integers and $a \neq 0$, we say that a **divides** b if there is an integer c such that $b = ac$. We write $a \mid b$ to say that a divides b , and $a \nmid b$ to say that a does not divide b .

Mathematically: $a \mid b \leftrightarrow \exists c \in \mathbb{Z} (b = ac)$

Note: If $a \mid b$, then

- a is called a **factor** of b
- b is called a **multiple** of a

We’ve been using the notion of divisibility all along!

- $E = \{x \mid x = 2k \wedge k \in \mathbb{Z}\}$



Division examples

Examples:

- Does $4 \mid 16$?
- Does $3 \mid 11$?
- Does $7 \mid 42$?

Question: Let n and d be two positive integers. How many positive integers not exceeding n are divisible by d ?

Answer: We want to count the number of integers of the form dk that are less than n . That is, we want to know the number of integers k with $0 \leq dk \leq n$, or $0 \leq k \leq n/d$. Therefore, there are $\lfloor n/d \rfloor$ positive integers not exceeding n that are divisible by d .



Important properties of divisibility

Property 1: If $a \mid b$ and $a \mid c$, then $a \mid (b + c)$

Proof: If $a \mid b$ and $a \mid c$, then there exist integers j and k such that $b = aj$ and $c = ak$. Hence, $b + c = aj + ak = a(j + k)$. Thus, $a \mid (b + c)$.

Property 2: If $a \mid b$, then $a \mid bc$ for all integers c .

Proof: If $a \mid b$, then this is some integer j such that $b = aj$. Multiplying both sides by c gives us $bc = ajc$, so by definition, $a \mid bc$.



One more property

Property 3: If $a \mid b$ and $b \mid c$, then $a \mid c$.

Proof: If $a \mid b$ and $b \mid c$, then there exist integers j and k such that $b = aj$ and $c = bk$. By substitution, we have that $c = ajk$, so $a \mid c$.



Division algorithm

Theorem: Let a be an integer and let d be a positive integer. There are unique integers q and r , with $0 \leq r < d$, such that $a = dq + r$.

For historical reasons, the above theorem is called **the division algorithm**, even though it isn't an algorithm!

Terminology: Given $a = dq + r$

- d is called the **divisor**
- q is called the **quotient**
- r is called the **remainder**
- $q = a \operatorname{div} d$
- $r = a \operatorname{mod} d$

Examples



Question: What are the quotient and remainder when 123 is divided by 23?

Answer: We have that $123 = 23 \times 5 + 8$. So the quotient is $123 \text{ div } 23 = 5$, and the remainder is $123 \bmod 23 = 8$.

Question: What are the quotient and remainder when -11 is divided by 3?

Answer: Since $-11 = 3 \times -4 + 1$, we have that the quotient is -11 and the remainder is 1.

Recall that since the remainder **must** be positive, $3 \times -3 - 2$ is not a valid use of the division theorem!

Many programming languages use the **div** and **mod** operations



For example, in Java, C, and C++

- / corresponds to **div** when used on integer arguments
- % corresponds to **mod**

```

public static void main(String[] args)
{
    int x = 2;
    int y = 5;
    float z = 2.0;

    System.out.println(y/x);
    System.out.println(y%x);
    System.out.println(y/z);
}

```

Prints out 1

→

Prints out 2, not 2.5!

Prints out 2.5

This can be a source of **many** errors, so be careful in your future classes!

Group work!



Problem 1: Does

1. $12 \mid 144$
2. $4 \mid 67$
3. $9 \mid 81$

Problem 2: What are the quotient and remainder when

1. 64 is divided by 8
2. 42 is divided by 11
3. 23 is divided by 7

Sometimes, we care only about the remainder of an integer after it is divided by some other integer



Example: What time will it be 22 hours from now?



Answer: If it is 6am now, it will be $(6 + 22) \bmod 24 = 28 \bmod 24 = 4$ am in 22 hours.

Since remainders can be so important, they have their own special notation!



Definition: If a and b are integers and m is a positive integer, we say that a is congruent to b modulo m if $m \mid (a - b)$. We write this as $a \equiv b \pmod{m}$.

Note: $a \equiv b \pmod{m}$ iff $a \bmod m = b \bmod m$.

Examples:

- Is 17 congruent to 5 modulo 6?
- Is 24 congruent to 14 modulo 6?

Properties of congruencies



Theorem: Let m be a positive integer. The integers a and b are congruent modulo m iff there is an integer k such that $a = b + km$.

Theorem: Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then

- $(a + c) \equiv (b + d) \pmod{m}$
- $ac \equiv bd \pmod{m}$

Congruencies have many applications within computer science



Today we'll look at two of the book's three:


- 1. Hash functions
- 2. Cryptography

Hash functions allow us to quickly and efficiently locate data



Problem: Given a large collection of records, how can we find the one we want quickly?

Solution: Apply a **hash function** that determines the storage location of the record based on the record's ID. A common hash function is $h(k) = k \bmod n$, where n is the number of available storage locations.



Hash functions are not one-to-one, so we must expect occasional collisions

Solution 1: Use next available location

Memory:

0	1	2	3	4	5	6	7
---	---	---	---	---	---	---	---

$42 \bmod 8 = 2$ $10 \bmod 8 = 2$

ID: 42
...
...

ID: 10
...
...

The field of cryptography makes heavy use of number theory and congruencies

Cryptography is the study of **secret messages**

Uses of cryptography:

- Protecting medical records
- Storing and transmitting military secrets
- Secure web browsing
- ...



Congruencies are used in cryptosystems from antiquity, as well as in modern-day algorithms

Since modern algorithms require quite a bit of sophistication to discuss, we'll examine an ancient cryptosystem

The Caesar cipher is based on congruencies



To encode a message using the Caesar cipher:

- Choose a shift index s
- Convert each letter A-Z into a number 0-25
- Compute $f(p) = p + s \bmod 26$

Example: Let $s = 9$. Encode “ATTACK”.

- ATTACK = 0 19 19 0 2 10
- $f(0) = 9, f(19) = 2, f(2) = 11, f(10) = 19$
- Encrypted message: 9 2 2 9 11 19 = JCCJLT

Decryption involves using the inverse function



That is, $f^{-1}(p) = p - s \bmod 26$

Example: Assume that $s = 3$. Decrypt the message “UHWUHDW”.

- UHWUHDW = 20 7 22 20 7 3 22
- $f^{-1}(20) = 17, f^{-1}(7) = 4, f^{-1}(22) = 19, f^{-1}(3) = 0$
- Decrypted result: 17 4 19 17 4 0 19 = RETREAT



Group work!

Problem 1:

1. Is 4 congruent to 8 mod 3?
2. Is 45 congruent to 12 mod 9?
3. Is 21 congruent to 28 mod 7?

Problem 2: The message “ROVY” was encrypted with the Caesar cipher using $s = 10$. Decrypt it.



Final thoughts

- Number theory is the study of integers and their properties
- Divisibility, modular arithmetic, and congruency are used throughout computer science
- Next time:
 - Prime numbers, GCDs, integer representation (Section 3.5)