

# CS 2740 Knowledge Representation

## Lecture 16

### Modeling uncertainty

**Milos Hauskrecht**

[milos@cs.pitt.edu](mailto:milos@cs.pitt.edu)

5329 Sennott Square

### KB systems. Medical example.

We want to build a KB system for the **diagnosis of pneumonia**.

#### Problem description:

- **Disease:** pneumonia
- **Patient symptoms (findings, lab tests):**
  - Fever, Cough, Paleness, WBC (white blood cells) count, Chest pain, etc.


#### Representation of a patient case:

- Statements that hold (are true) for the patient.
  - E.g:       Fever =*True*
  - Cough =*False*
  - WBCcount=*High*

**Diagnostic task:** we want to decide whether the patient suffers from the pneumonia or not given the symptoms

## Uncertainty

To make diagnostic inference possible we need to represent knowledge (axioms) that relate symptoms and diagnosis



**Problem:** disease/symptoms relations are not deterministic

- They are **uncertain (or stochastic)** and vary from patient to patient

## Uncertainty

**Two types of uncertainty:**

- **Disease → Symptoms uncertainty**

- A patient suffering from pneumonia may not have fever all the times, may or may not have a cough, white blood cell test can be in a normal range.

- **Symptoms → Disease uncertainty**

- High fever is typical for many diseases (e.g. bacterial diseases) and does not point specifically to pneumonia
- Fever, cough, paleness, high WBC count combined do not always point to pneumonia

## Uncertainty

### Why are relations uncertain?

- **Observability**

- It is impossible to observe all relevant components of the world
- Observable components behave stochastically even if the underlying world is deterministic

- **Efficiency, capacity limits**

- It is often impossible to enumerate and model all components of the world and their relations
- abstractions can become stochastic


### **Humans can reason with uncertainty !!!**

- Can computer systems do the same?

## Modeling the uncertainty.

### **Key challenges:**

- How to represent the relations in the presence of uncertainty?
- How to manipulate such knowledge to make inferences?
  - **Humans can reason with uncertainty.**



## Methods for representing uncertainty

### Extensions of the propositional and first-order logic

- Use, uncertain, imprecise statements (relations)

#### Example: Propositional logic with certainty factors

Very popular in 70-80s in knowledge-based systems (MYCIN)

- **Facts (propositional statements)** are assigned a **certainty value** reflecting the belief in that the statement is satisfied:

$$CF(Pneumonia = True) = 0.7$$

- **Knowledge:** typically in terms of **modular rules**

<b>If</b>	1. The patient has cough, and 2. The patient has a high WBC count, and 3. The patient has fever
<b>Then</b>	<b>with certainty 0.7</b> the patient has pneumonia

## Certainty factors

### Problem 1:

- Chaining of multiple inference rules (propagation of uncertainty)

### Solution:

- **Rules** incorporate tests on the **certainty values**

$$(A \text{ in } [0.5,1]) \wedge (B \text{ in } [0.7,1]) \rightarrow C \text{ with } CF = 0.8$$

### Problem 2:

- Combinations of rules **with the same conclusion**

$$(A \text{ in } [0.5,1]) \wedge (B \text{ in } [0.7,1]) \rightarrow C \text{ with } CF = 0.8$$

$$(E \text{ in } [0.8,1]) \wedge (D \text{ in } [0.9,1]) \rightarrow C \text{ with } CF = 0.9$$

- What is the resulting  $CF(C)$  ?

## Certainty factors

- Combination of multiple rules

$(A \text{ in } [0.5,1]) \wedge (B \text{ in } [0.7,1]) \rightarrow C$  with  $CF = 0.8$

$(E \text{ in } [0.8,1]) \wedge (D \text{ in } [0.9,1]) \rightarrow C$  with  $CF = 0.9$

- Three possible solutions

$$CF(C) = \max[0.9; 0.8] = 0.9$$

$$CF(C) = 0.9 * 0.8 = 0.72$$

$$CF(C) = 0.9 + 0.8 - 0.9 * 0.8 = 0.98$$

} ?

**Problems:**

- Which solution to choose?
- All three methods break down after a sequence of inference rules

## Methods for representing uncertainty

### Probability theory

- A well defined theory for modeling and reasoning in the presence of uncertainty
- A natural choice to replace certainty factors

### Facts (propositional statements)

- Are represented via **random variables** with two or more values

**Example:** *Pneumonia* is a random variable

**values:** *True* and *False*

- Each value can be achieved **with some probability:**

$$P(Pneumonia = True) = 0.001$$

$$P(WBCcount = high) = 0.005$$


## Probability theory

- Well-defined theory for representing and manipulating statements with uncertainty

- **Axioms of probability:**

For any two propositions A, B.

1.  $0 \leq P(A) \leq 1$
2.  $P(\text{True}) = 1$  and  $P(\text{False}) = 0$
3.  $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$



## Modeling uncertainty with probabilities

### Probabilistic extension of propositional logic.

- **Propositions:**

- statements about the world
- Represented by the assignment of values to **random variables**

- **Random variables:**

- ! – **Boolean**      *Pneumonia* is either *True, False*  

Random variable	Values
-----------------	--------
- ! – **Multi-valued**      *Pain* is one of  $\{Nopain, Mild, Moderate, Severe\}$   

Random variable	Values
-----------------	--------
- **Continuous**      *HeartRate* is a value in  $<0; 250>$   

Random variable	Values
-----------------	--------

## Probabilities

### Measure the degree of our belief in propositions

$$P(\textit{Pneumonia} = \textit{True}) = 0.001$$

$$P(\textit{Pneumonia} = \textit{False}) = 0.999$$

$$P(\textit{WBCcount} = \textit{high}) = 0.005$$

### Probability distribution

- Defines probabilities **for all possible value assignments to a random variable**
- Values are mutually exclusive

$$P(\textit{Pneumonia} = \textit{True}) = 0.001$$

$$P(\textit{Pneumonia} = \textit{False}) = 0.999$$

<i>Pneumonia</i>	<b>P</b> ( <i>Pneumonia</i> )
<i>True</i>	0.001
<i>False</i>	0.999

## Probability distribution

Defines probability for **all possible value assignments**

### Example 1:

$$P(\textit{Pneumonia} = \textit{True}) = 0.001$$

$$P(\textit{Pneumonia} = \textit{False}) = 0.999$$

<i>Pneumonia</i>	<b>P</b> ( <i>Pneumonia</i> )
<i>True</i>	0.001
<i>False</i>	0.999

$$P(\textit{Pneumonia} = \textit{True}) + P(\textit{Pneumonia} = \textit{False}) = 1$$

**Probabilities sum to 1 !!!**

### Example 2:

$$P(\textit{WBCcount} = \textit{high}) = 0.005$$

$$P(\textit{WBCcount} = \textit{normal}) = 0.993$$

$$P(\textit{WBCcount} = \textit{low}) = 0.002$$

<i>WBCcount</i>	<b>P</b> ( <i>WBCcount</i> )
<i>high</i>	0.005
<i>normal</i>	0.993
<i>low</i>	0.002

## Joint probability distribution

### Joint probability distribution (for a set variables)

- Defines probabilities for **all possible assignments of values to variables in the set**

**Example:** variables *Pneumonia* and *WBCcount*

$P(\text{pneumonia}, \text{WBCcount})$

Is represented by  $2 \times 3$  matrix

		<i>WBCcount</i>		
		<i>high</i>	<i>normal</i>	<i>low</i>
<i>Pneumonia</i>	<i>True</i>	0.0008	0.0001	0.0001
	<i>False</i>	0.0042	0.9929	0.0019

## Joint probabilities

### Marginalization

- reduces the dimension of the joint distribution
- Sums variables out

$P(\text{pneumonia}, \text{WBCcount})$   $2 \times 3$  matrix

		<i>WBCcount</i>			
		<i>high</i>	<i>normal</i>	<i>low</i>	
<i>Pneumonia</i>	<i>True</i>	0.0008	0.0001	0.0001	0.001
	<i>False</i>	0.0042	0.9929	0.0019	
		0.005	0.993	0.002	

$P(\text{WBCcount})$

**Marginalization** (here summing of columns or rows)



## Full joint distribution

- the joint distribution for all variables in the problem
  - It defines the complete probability model for the problem

**Example:** pneumonia diagnosis

**Variables:** *Pneumonia, Fever, Paleness, WBCcount, Cough*

Full joint defines the probability for all possible assignments of values to *Pneumonia, Fever, Paleness, WBCcount, Cough*

$P(\text{Pneumonia} = T, \text{WBCcount} = \text{High}, \text{Fever} = T, \text{Cough} = T, \text{Paleness} = T)$

$P(\text{Pneumonia} = T, \text{WBCcount} = \text{High}, \text{Fever} = T, \text{Cough} = T, \text{Paleness} = F)$

$P(\text{Pneumonia} = T, \text{WBCcount} = \text{High}, \text{Fever} = T, \text{Cough} = F, \text{Paleness} = T)$

... etc

## Conditional probabilities

### Conditional probability distribution

- Defines probabilities for all possible assignments, given a fixed assignment to some other variable values

$P(\text{Pneumonia} = \text{true} \mid \text{WBCcount} = \text{high})$

$P(\text{Pneumonia} \mid \text{WBCcount})$  3 element vector of 2 elements

		<i>WBCcount</i>		
		<i>high</i>	<i>normal</i>	<i>low</i>
<i>Pneumonia</i>	<i>True</i>	0.08	0.0001	0.0001
	<i>False</i>	0.92	0.9999	0.9999
		1.0	1.0	1.0

$P(\text{Pneumonia} = \text{true} \mid \text{WBCcount} = \text{high})$

+  $P(\text{Pneumonia} = \text{false} \mid \text{WBCcount} = \text{high})$

## Conditional probabilities

### Conditional probability

- Is defined in terms of the joint probability:

$$P(A|B) = \frac{P(A,B)}{P(B)} \text{ s.t. } P(B) \neq 0$$

- Example:**

$$P(\text{pneumonia} = \text{true} | \text{WBCcount} = \text{high}) = \frac{P(\text{pneumonia} = \text{true}, \text{WBCcount} = \text{high})}{P(\text{WBCcount} = \text{high})}$$

$$P(\text{pneumonia} = \text{false} | \text{WBCcount} = \text{high}) = \frac{P(\text{pneumonia} = \text{false}, \text{WBCcount} = \text{high})}{P(\text{WBCcount} = \text{high})}$$

## Conditional probabilities

- Conditional probability distribution.**

$$P(A|B) = \frac{P(A,B)}{P(B)} \text{ s.t. } P(B) \neq 0$$

- Product rule.** Joint probability can be expressed in terms of conditional probabilities

$$P(A,B) = P(A|B)P(B)$$

- Chain rule.** Any joint probability can be expressed as a product of conditionals

$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_n | X_1, \dots, X_{n-1})P(X_1, \dots, X_{n-1}) \\ &= P(X_n | X_1, \dots, X_{n-1})P(X_{n-1} | X_1, \dots, X_{n-2})P(X_1, \dots, X_{n-2}) \\ &= \prod_{i=1}^n P(X_i | X_1, \dots, X_{i-1}) \end{aligned}$$

## Bayes rule

Conditional probability.

$$P(A|B) = \frac{P(A,B)}{P(B)} \quad \text{where} \quad P(A,B) = P(B|A)P(A)$$

Bayes rule:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

When is it useful?

- When we are interested in computing the diagnostic query from the causal probability

$$P(\text{cause} | \text{effect}) = \frac{P(\text{effect} | \text{cause})P(\text{cause})}{P(\text{effect})}$$

- **Reason:** It is often easier to assess causal probability
  - E.g. Probability of pneumonia causing fever  
vs. probability of pneumonia given fever

## Bayes rule

Assume a variable A with multiple values  $a_1, a_2, \dots, a_k$

Bayes rule can be rewritten as:


$$\begin{aligned} P(A = a_j | B = b) &= \frac{P(B = b | A = a_j)P(A = a_j)}{P(B = b)} \\ &= \frac{P(B = b | A = a_j)P(A = a_j)}{\sum_{i=1}^k P(B = b | A = a_i)P(A = a_i)} \end{aligned}$$

Used in practice when we want to compute:

$$P(A | B = b) \quad \text{for all values of} \quad a_1, a_2, \dots, a_k$$

## Bayes Rule in a simple diagnostic inference

- **Device** (equipment) operating *normally* or *malfunctioning*.
  - Operation of the device sensed indirectly via a sensor
- **Sensor reading** is either *high* or *low*



**P(Device status)**

<b>normal</b>	<b>malfunctioning</b>
0.9	0.1

**P(Sensor reading | Device status)**

Device\Sensor	<b>high</b>	<b>low</b>
normal	0.1	0.9
malfunctioning	0.6	0.4

## Bayes Rule in a simple diagnostic inference.

- **Diagnostic inference:** compute the probability of device operating normally or malfunctioning given a sensor reading

$$P(\text{Device status} \mid \text{Sensor reading} = \text{high}) = ?$$

$$= \begin{pmatrix} P(\text{Device status} = \text{normal} \mid \text{Sensor reading} = \text{high}) \\ P(\text{Device status} = \text{malfunctioning} \mid \text{Sensor reading} = \text{high}) \end{pmatrix}$$

- Note that typically the opposite conditional probabilities are given to us: they are much easier to estimate
- **Solution:** apply **Bayes rule** to reverse the conditioning variables

## Probabilistic inference

Various inference tasks:

- **Diagnostic task. (from effect to cause)**

$$\mathbf{P}(Pneumonia \mid Fever = T)$$

- **Prediction task. (from cause to effect)**

$$\mathbf{P}(Fever \mid Pneumonia = T)$$

- **Other probabilistic queries** (queries on joint distributions).

$$\mathbf{P}(Fever)$$

$$\mathbf{P}(Fever, ChestPain)$$