

# CS 441 Discrete Mathematics for CS

## Lecture 35

### Relations

**Milos Hauskrecht**

[milos@cs.pitt.edu](mailto:milos@cs.pitt.edu)

5329 Sennott Square

### Course administration

- **Homework assignment 11**
  - due on Friday, April 21, 2006
- **Final exam**
  - Thursday, April 27, 2006
  - At 12:00-1:50pm
  - The same room as lectures

**Course web page:**

<http://www.cs.pitt.edu/~milos/courses/cs441/>

## Cartesian product (review)

- Let  $A = \{a_1, a_2, \dots, a_k\}$  and  $B = \{b_1, b_2, \dots, b_m\}$ .
- **The Cartesian product**  $A \times B$  is defined by a set of pairs  $\{(a_1, b_1), (a_1, b_2), \dots, (a_1, b_m), \dots, (a_k, b_m)\}$ .

### Example:

Let  $A = \{a, b, c\}$  and  $B = \{1, 2, 3\}$ . What is  $A \times B$ ?

## Cartesian product (review)

- Let  $A = \{a_1, a_2, \dots, a_k\}$  and  $B = \{b_1, b_2, \dots, b_m\}$ .
- **The Cartesian product**  $A \times B$  is defined by a set of pairs  $\{(a_1, b_1), (a_1, b_2), \dots, (a_1, b_m), \dots, (a_k, b_m)\}$ .

### Example:

Let  $A = \{a, b, c\}$  and  $B = \{1, 2, 3\}$ . What is  $A \times B$ ?

$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$

## Binary relation

**Definition:** Let  $A$  and  $B$  be sets. A **binary relation from  $A$  to  $B$**  is a subset of a Cartesian product  $A \times B$ .

**Example:** Let  $A = \{a, b, c\}$  and  $B = \{1, 2, 3\}$ .

- $R = \{(a, 1), (b, 2), (c, 2)\}$  is an example of a relation from  $A$  to  $B$ .

## Binary relation

**Definition:** Let  $A$  and  $B$  be sets. A **binary relation from  $A$  to  $B$**  is a subset of a Cartesian product  $A \times B$ .

**Example:** Let  $A = \{a, b, c\}$  and  $B = \{1, 2, 3\}$ .

- $R = \{(a, 1), (b, 2), (c, 2)\}$  is an example of a relation from  $A$  to  $B$ .
- Another example of a relation from  $A$  to  $B$ ?


## Representing binary relations

- We can graphically represent a binary relation  $R$  as follows:
  - if  $a R b$  then draw an arrow from  $a$  to  $b$ .

$$a \rightarrow b$$

### Example:

- Let  $A = \{0, 1, 2\}$ ,  $B = \{u, v\}$  and  $R = \{ (0, u), (0, v), (1, v), (2, u) \}$
- Note:  $R \subseteq A \times B$ .
- Graph:**



## Representing binary relations

- We can represent a binary relation  $R$  by a **table** showing (marking) the ordered pairs of  $R$ .

### Example:


- Let  $A = \{0, 1, 2\}$ ,  $B = \{u, v\}$  and  $R = \{ (0, u), (0, v), (1, v), (2, u) \}$
- Table:**

R	u	v	or	R	u	v
0	x	x		0	1	1
1		x		1	0	1
2	x			2	1	0


## Relations and functions

- Relations represent **one to many relationships** between elements in A and B.

- Example:**



- What is the difference between a **relation** and a **function from A to B**? A function on sets A,B  $A \rightarrow B$  assigns to each element in the domain set A exactly one element from B. So it is a **special relation**.



## Relation on the set

**Definition:** A relation on the set A is a relation from A to itself.

### Example 1:

- Let  $A = \{1,2,3,4\}$  and
- $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  is defined as:
  - $R_{\text{fun}} = \{(1,2),(2,2),(3,3)\}$ .

## Relation on the set

**Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example 1:**

- Let  $A = \{1,2,3,4\}$  and  $R_{\text{div}} = \{(a,b) \mid a \text{ divides } b\}$
- What does  $R_{\text{div}}$  consist of?
- $R_{\text{div}} = \{ \dots$

## Relation on the set

**Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example 1:**

- Let  $A = \{1,2,3,4\}$  and  $R_{\text{div}} = \{(a,b) \mid a \text{ divides } b\}$
- What does  $R_{\text{div}}$  consist of?
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$

R	1	2	3	4
1	x	x	x	x
2		x		x
3			x	
4				x

## Relation on the set

### Example 2:

- Let  $A = \{1,2,3,4\}$ .
- Define  $a R_{\neq} b$  if and only if  $a \neq b$ .

$R_{\neq} = \{ \dots$

## Relation on the set

### Example 2:

- Let  $A = \{1,2,3,4\}$ .
- Define  $a R_{\neq} b$  if and only if  $a \neq b$ .

$R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$

•	<table><tr><th>R</th><th>1</th><th>2</th><th>3</th><th>4</th></tr><tr><td>1</td><td></td><td>x</td><td>x</td><td>x</td></tr><tr><td>2</td><td>x</td><td></td><td>x</td><td>x</td></tr><tr><td>3</td><td>x</td><td>x</td><td></td><td>x</td></tr><tr><td>4</td><td>x</td><td>x</td><td>x</td><td></td></tr></table>	R	1	2	3	4	1		x	x	x	2	x		x	x	3	x	x		x	4	x	x	x	
R	1	2	3	4																						
1		x	x	x																						
2	x		x	x																						
3	x	x		x																						
4	x	x	x																							

## Relation on the set

**Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example 3:**

- Let  $A = \{1,2,3,4\}$  and
- $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  is defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .

## Properties of relations

**Definition (reflexive relation) :** A relation  $R$  on a set  $A$  is called **reflexive** if  $(a,a) \in R$  for every element  $a \in A$ .

**Example 1:**

- Assume relation  $R_{\text{div}} = \{(a,b) \mid a \mid b\}$  on  $A = \{1,2,3,4\}$
- **Is  $R_{\text{div}}$  reflexive?**
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Answer: ?**



## Properties of relations

**Definition (reflexive relation)** : A relation  $R$  on a set  $A$  is called **reflexive** if  $(a,a) \in R$  for every element  $a \in A$ .

### Example 1:

- Assume relation  $R_{\text{div}} = \{(a,b) \mid a \mid b\}$  on  $A = \{1,2,3,4\}$
- **Is  $R_{\text{div}}$  reflexive?**
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Answer:** Yes.  $(1,1), (2,2), (3,3),$  and  $(4,4) \in A$ .

## Reflexive relation

### Reflexive relation

- $R_{\text{div}} = \{(a,b) \mid a \mid b\}$  on  $A = \{1,2,3,4\}$
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$

$$\text{MR}_{\text{div}} = \begin{array}{cccc} & 1 & 2 & 3 & 4 \\ \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \end{array} & \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \end{array} & \begin{array}{c} 1 \\ 1 \\ 0 \\ 0 \end{array} & \begin{array}{c} 1 \\ 0 \\ 1 \\ 0 \end{array} & \begin{array}{c} 1 \\ 1 \\ 0 \\ 1 \end{array} \end{array}$$

- **A relation  $R$  is reflexive** if and only if  $MR$  has 1 in every position on its main diagonal.

## Properties of relations

**Definition (reflexive relation)** : A relation  $R$  on a set  $A$  is called **reflexive** if  $(a,a) \in R$  for every element  $a \in A$ .

### Example 2:

- Relation  $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .
- **Is  $R_{\text{fun}}$  reflexive?**

## Properties of relations

**Definition (reflexive relation)** : A relation  $R$  on a set  $A$  is called **reflexive** if  $(a,a) \in R$  for every element  $a \in A$ .

### Example 2:

- Relation  $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .
- **Is  $R_{\text{fun}}$  reflexive?**
- **No.** It is not reflexive since  $(1,1) \notin R_{\text{fun}}$ .

## Properties of relations

**Definition (irreflexive relation):** A relation  $R$  on a set  $A$  is called **irreflexive** if  $(a,a) \notin R$  for every  $a \in A$ .

### Example 1:

- Assume relation  $R_{\neq}$  on  $A = \{1,2,3,4\}$ , such that  $a R_{\neq} b$  if and only if  $a \neq b$ .
- Is  $R_{\neq}$  irreflexive?
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Answer:**

## Properties of relations

**Definition (irreflexive relation):** A relation  $R$  on a set  $A$  is called **irreflexive** if  $(a,a) \notin R$  for every  $a \in A$ .

### Example 1:

- Assume relation  $R_{\neq}$  on  $A = \{1,2,3,4\}$ , such that  $a R_{\neq} b$  if and only if  $a \neq b$ .
- Is  $R_{\neq}$  irreflexive?
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Answer:** Yes. Because  $(1,1), (2,2), (3,3)$  and  $(4,4) \notin R_{\neq}$

## Irreflexive relation

### Irreflexive relation

- $R_{\neq}$  on  $A = \{1, 2, 3, 4\}$ , such that  $\mathbf{a} R_{\neq} \mathbf{b}$  if and only if  $a \neq b$ .
- $R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}$

		0	1	1	1
		1	0	1	1
MR	=	1	1	0	1
		1	1	1	0

- **A relation  $R$  is irreflexive** if and only if  $MR$  has 0 in every position on its main diagonal.

## Properties of relations

**Definition (irreflexive relation):** A relation  $R$  on a set  $A$  is called **irreflexive** if  $(a, a) \notin R$  for every  $a \in A$ .

### Example 2:

- $R_{\text{fun}}$  on  $A = \{1, 2, 3, 4\}$  defined as:
  - $R_{\text{fun}} = \{(1, 2), (2, 2), (3, 3)\}$ .
- **Is  $R_{\text{fun}}$  irreflexive?**
- **Answer:**

## Properties of relations

**Definition (irreflexive relation):** A relation  $R$  on a set  $A$  is called **irreflexive** if  $(a,a) \notin R$  for every  $a \in A$ .

### Example 2:

- $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .
- **Is  $R_{\text{fun}}$  irreflexive?**
- **Answer: No.** Because  $(2,2)$  and  $(3,3) \in R_{\text{fun}}$

## Properties of relations

**Definition (symmetric relation):** A relation  $R$  on a set  $A$  is called **symmetric** if

$$\forall a, b \in A \quad (a,b) \in R \rightarrow (b,a) \in R.$$

### Example 1:

- $R_{\text{div}} = \{(a,b) \mid a \mid b\}$  on  $A = \{1,2,3,4\}$
- **Is  $R_{\text{div}}$  symmetric?**
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Answer:**

## Properties of relations

**Definition (symmetric relation):** A relation  $R$  on a set  $A$  is called **symmetric** if

$$\forall a, b \in A \quad (a, b) \in R \rightarrow (b, a) \in R.$$

**Example 1:**

- $R_{\text{div}} = \{(a, b) \mid a \mid b\}$  on  $A = \{1, 2, 3, 4\}$
- **Is  $R_{\text{div}}$  symmetric?**
- $R_{\text{div}} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}$
- **Answer: No.** It is not symmetric since  $(1, 2) \in R$  but  $(2, 1) \notin R$ .

## Properties of relations

**Definition (symmetric relation):** A relation  $R$  on a set  $A$  is called **symmetric** if

$$\forall a, b \in A \quad (a, b) \in R \rightarrow (b, a) \in R.$$

**Example 2:**

- $R_{\neq}$  on  $A = \{1, 2, 3, 4\}$ , such that  $a R_{\neq} b$  if and only if  $a \neq b$ .
- **Is  $R_{\neq}$  symmetric?**
- $R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}$
- **Answer:**

## Properties of relations

**Definition (symmetric relation):** A relation  $R$  on a set  $A$  is called **symmetric** if

$$\forall a, b \in A \quad (a, b) \in R \rightarrow (b, a) \in R.$$

**Example 2:**

- $R_{\neq}$  on  $A = \{1, 2, 3, 4\}$ , such that  **$a R_{\neq} b$**  if and only if  $a \neq b$ .
- **Is  $R_{\neq}$  symmetric ?**
- $R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}$
- **Answer: Yes.** If  $(a, b) \in R_{\neq} \rightarrow (b, a) \in R_{\neq}$

## Symmetric relation

**Symmetric relation:**

- $R_{\neq}$  on  $A = \{1, 2, 3, 4\}$ , such that  **$a R_{\neq} b$**  if and only if  $a \neq b$ .
- $R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}$

$$\text{MR} = \begin{array}{cccc} & 0 & 1 & 1 & 1 \\ & 1 & 0 & 1 & 1 \\ & 1 & 1 & 0 & 1 \\ & 1 & 1 & 1 & 0 \end{array}$$

- **A relation  $R$  is symmetric** if and only if  $m_{ij} = m_{ji}$  for all  $i, j$ .

## Properties of relations

- **Definition (antisymmetric relation):** A relation on a set  $A$  is called **antisymmetric** if
  - $[(a,b) \in R \text{ and } (b,a) \in R] \rightarrow a = b$  where  $a, b \in A$ .

### Example 1:

- Relation  $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .
- **Is  $R_{\text{fun}}$  antisymmetric?**
- **Answer:**

## Properties of relations

- **Definition (antisymmetric relation):** A relation on a set  $A$  is called **antisymmetric** if
  - $[(a,b) \in R \text{ and } (b,a) \in R] \rightarrow a = b$  where  $a, b \in A$ .

### Example 1:

- Relation  $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .
- **Is  $R_{\text{fun}}$  antisymmetric?**
- **Answer: Yes.** It is antisymmetric



## Antisymmetric relations

### Antisymmetric relation

- relation  $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$

$$MR_{\text{fun}} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$

- A relation is **antisymmetric** if and only if  $m_{ij} = 1 \rightarrow m_{ji} = 0$  for  $i \neq j$ .

## Properties of relations

**Definition (antisymmetric relation):** A relation on a set  $A$  is called **antisymmetric** if

- $[(a,b) \in R \text{ and } (b,a) \in R] \rightarrow a = b$  where  $a, b \in A$ .

### Example 2:

- $R_{\neq}$  on  $A = \{1,2,3,4\}$ , such that  $a R_{\neq} b$  if and only if  $a \neq b$ .
- Is  $R_{\neq}$  antisymmetric ?
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- Answer:**

## Properties of relations

**Definition (antisymmetric relation):** A relation on a set  $A$  is called **antisymmetric** if

- $[(a,b) \in R \text{ and } (b,a) \in R] \rightarrow a = b$  where  $a, b \in A$ .

**Example 2:**

- $R_{\neq}$  on  $A = \{1,2,3,4\}$ , such that  $a R_{\neq} b$  if and only if  $a \neq b$ .
- **Is  $R_{\neq}$  antisymmetric ?**
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Answer: No.** It is not antisymmetric since  $(1,2) \in R$  and  $(2,1) \in R$  but  $1 \neq 2$ .

## Properties of relations

**Definition (transitive relation):** A relation  $R$  on a set  $A$  is called **transitive** if

- $[(a,b) \in R \text{ and } (b,c) \in R] \rightarrow (a,c) \in R$  for all  $a, b, c \in A$ .

• **Example 1:**

- $R_{\text{div}} = \{(a,b) \mid a \mid b\}$  on  $A = \{1,2,3,4\}$
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Is  $R_{\text{div}}$  transitive?**
- **Answer:**

## Properties of relations

**Definition (transitive relation):** A relation  $R$  on a set  $A$  is called **transitive** if

- $[(a,b) \in R \text{ and } (b,c) \in R] \rightarrow (a,c) \in R$  for all  $a, b, c \in A$ .

- **Example 1:**

- $R_{\text{div}} = \{(a,b) \mid a \mid b\}$  on  $A = \{1,2,3,4\}$
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Is  $R_{\text{div}}$  transitive?**
- **Answer: Yes.**

## Properties of relations

**Definition (transitive relation):** A relation  $R$  on a set  $A$  is called **transitive** if

- $[(a,b) \in R \text{ and } (b,c) \in R] \rightarrow (a,c) \in R$  for all  $a, b, c \in A$ .

- **Example 2:**

- $R_{\neq}$  on  $A = \{1,2,3,4\}$ , such that  $a R_{\neq} b$  if and only if  $a \neq b$ .
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Is  $R_{\neq}$  transitive ?**
- **Answer:**

## Properties of relations

**Definition (transitive relation):** A relation  $R$  on a set  $A$  is called **transitive** if

- $[(a,b) \in R \text{ and } (b,c) \in R] \rightarrow (a,c) \in R$  for all  $a, b, c \in A$ .
- **Example 2:**
- $R_{\neq}$  on  $A = \{1,2,3,4\}$ , such that  $\mathbf{a R_{\neq} b}$  if and only if  $a \neq b$ .
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Is  $R_{\neq}$  transitive?**
- **Answer: No.** It is not transitive since  $(1,2) \in R$  and  $(2,1) \in R$  but  $(1,1)$  is not an element of  $R$ .

## Properties of relations

**Definition (transitive relation):** A relation  $R$  on a set  $A$  is called **transitive** if

- $[(a,b) \in R \text{ and } (b,c) \in R] \rightarrow (a,c) \in R$  for all  $a, b, c \in A$ .
- **Example 3:**
- Relation  $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:
  - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .
- **Is  $R_{\text{fun}}$  transitive?**
- **Answer:**

## Properties of relations

**Definition (transitive relation):** A relation  $R$  on a set  $A$  is called **transitive** if

- $[(a,b) \in R \text{ and } (b,c) \in R] \rightarrow (a,c) \in R$  for all  $a, b, c \in A$ .

- **Example 3:**

- Relation  $R_{\text{fun}}$  on  $A = \{1,2,3,4\}$  defined as:

- $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$ .

- **Is  $R_{\text{fun}}$  transitive?**

- **Answer: Yes.** It is transitive.